

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification 6:		(11) International Publication Number: WO 98/11067	
C07D 211/26, 498/04 // (C07D 498/04, 265:00, 209:00)	A1	(43) International Publication Date: 19 March 1998 (19.03.98)	
 (21) International Application Number: PCT/EP (22) International Filing Date: 9 September 1997 ((30) Priority Data: 9618967.5 11 September 1996 (11.09.9) (71) Applicant (for all designated States except US): SMIT BEECHAM P.L.C. [GB/GB]; New Horizons Courford, Middlesex TW8 9EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): FEDOULOFF, [GB/GB]; SmithKline Beecham Pharmaceutica Frontiers Science Park South, Third Avenue, Harloc CM19 5AW (GB). GUEST, David, William [SmithKline Beecham Pharmaceuticals, New Science Park South, Third Avenue, Harlow, Esse 5AW (GB). SMITH, Gillian, Elizabeth [GB/GB]; line Beecham Pharmaceuticals, New Frontiers Science South, Third Avenue, Harlow, Essex CM19 5AW (74) Agent: CONNELL, Anthony, Christopher; Sn Beecham, Corporate Intellectual Property, Tw Horizons Court, Brentford, Middlesex TW8 9EP (6) 	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BF, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GF, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NC, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GF, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NI, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.		
	lmethyl:	HYLAMINE unine, which process comprises: i) the reaction of isonipecotamide and ollowed by ii) reduction with LiAlH ₄ , characterised in that the reactions	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	Fī	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
ВВ	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israe)	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of Americ
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	٧N	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwc
CI	Côte d'Ivoire	КP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		•
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

10

15

20

25

PREPARATION OF 1-BUTYL-4-PIPERIDINYLMETHYLAMINE

This invention relates to a new synthetic process to an intermediate which is useful for the preparation of compounds having pharmacological activity.

-1-

WO 93/03725, WO 93/05038, WO 93/08187, WO 93/16072, WO 93/18027, WO 93/18036, WO 94/07859, WO 94/08965, WO 94/08994, WO 94/08995, WO 94/08998, WO 94/17071 (SmithKline Beecham plc) describe compounds having 5-HT₄ receptor antagonist activity.

WO 93/18036, Example 3 describes N-[(1-nbutyl-4-piperidyl)methyl]-3,4dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide SB 207266, (the hydrochloride salt is SB 207266-A) which is being developed by SmithKline Beecham plc as the active ingredient in a medicament for treatment of irritable bowel syndrome.

WO 93/18036 describes a method of preparation of SB 207266-A from N-[(1nbutyl-4-piperidyl)methyl]indole-3-carboxamide (i.e. the compound corresponding to SB 207266, without the oxazino moiety), by reacting with N-chlorosuccinimide and 3bromo-1-propanol, followed by treatment with sodium carbonate. N-[(1-nbutyl-4piperidyl)methyl]indole-3-carboxamide is prepared by coupling 1-butyl-4piperidinylmethylamine with indole-3-carboxylic acid. The 1-butyl-4piperidinylmethylamine is prepared as in Description 7 of WO 93/05038 and Description 1 of WO 93/18036, in a three stage process from isonipecotamide and 1-bromobutane, by alkylation in ethanol, to give the N-butyl derivative of isonipecotamide which is dehydrated to the corresponding nitrile and then reduced with LiAlH4 in ether.

An alternative process for preparing 1-butyl-4-piperidinylmethylamine has now been discovered which involves the use of a common solvent, allowing the two stages to be run without isolation of the N-butyl derivative of isonipecotamide.

Accordingly, the present invention provides a process for the preparation of 1-butyl-4-piperidinylmethylamine, which process comprises:

- 30 i) the reaction of isonipecotamide and 1-bromobutane to give the N-butyl derivative of isonipecotamide; followed by
 - reduction with LiAlH4. ii)

characterised in that the reactions i) and ii) are carried out in toluene as solvent.

The advantages of this process as compared with that previously described are as follows:

- 1. Toluene does not contain any additives, whereas THF contains a stabiliser (dit-butylcresol) which can only be removed from 1-butyl-4-piperidinylmethylamine by fractional distillation.
 - 2. The overall process does not involve the preparation/isolation of the intermediate nitrile, and is therefore one step shorter.
- 3. The process does not involve the isolation of the N-butyl derivative of isonipecotamide.
 - 4. The process uses a single solvent and eliminates the use of ethanol, chloroform and THF.
 - 5. the special extractive work-up of the LiAlH₄ reaction reduces the usage of solvent and loss of product on solid alumina residues.
- 15 The following Examples illustrate the invention.

15

25

Example 1

4-Piperidinecarboxamide (*iso*-nipecotamide) and potassium carbonate (2 equivs.) were stirred in toluene and treated with 1-bromobutane (1 equiv.). The reaction mixture was heated at reflux (107-110°C) for 2 hours. After cooling to 80-85°C the mixture was washed with hot water followed by hot aqueous potassium carbonate solution. The resulting toluene solution of 1-butyl-*iso*-nipecotamide was dried by azeotropic distillation, maintaining the reaction volume by addition of fresh toluene.

The toluene solution was cooled to 0-5°C, under nitrogen. A solution of LiAlH₄.2THF in toluene (1.0 molar solution; 2.0 equivs.) was added over 1 hour, keeping the temperature <10°C. The mixture was allowed to warm to room temperature and was then heated to reflux for 1 hour. After cooling to 0-5°C, 32% w/w sodium hydroxide solution (1.5 equivs. wrt substrate) was added cautiously over 1 hour, keeping the temperature <10°C. The mixture was stirred for 30 minutes at ambient temperature and the precipitate filtered through celite, washing the bed thoroughly with toluene. The filtrate was evaporated *in vacuo* to give 1-butyl-4-piperidinylmethylamine as a pale yellow oil, containing ~13% by weight toluene, in 72% yield (after adjusting for toluene content).

20 Example 2

Alternatively the first part of the preparation may be carried out as follows:

4-Piperidinecarboxamide (*iso*-nipecotamide) and 5M aqueous potassium carbonate solution (2 equivs.) were stirred in toluene and treated with 1-bromobutane (1 equiv.). The reaction mixture was heated at reflux (107-110°C) for 2 hours. After cooling to 70-80°C the mixture was washed with hot water followed by hot aqueous potassium carbonate solution. The resulting toluene solution of 1-butyl-*iso*-nipecotamide was dried by azeotropic distillation, maintaining the reaction volume by addition of fresh toluene.

Example 3 A 3L vessel was purged with nitrogen and charged with *iso*nipecotamide (112.1g 0.87mol) and dry toluene (785ml). The suspension was warmed to 50°C and potassium carbonate (248g, 1.79mol) and butyl bromide (119.8g,

0.87mol) were added in one portion. The resulting mixture was heated at reflux under Dean-Stark conditions for three hours and then cooled to 65°C and quenched by addition of water (875ml). The aqueous phase was separated at about 80°C and the organic layer dried by azeotropic distillation of toluene (200ml). Fresh toluene (200ml) was added to maintain a constant volume.

The reaction mixture was cooled to about 5°C and treated, dropwise, with a solution of lithium aluminium hydride.2THF in toluene (500ml, 3.5M, 1.75mol). The mixture was stirred at ambient temperature for one hour and then at about 55°C for a further two hours. The reaction was then quenched by cautious addition of sodium hydroxide solution (1200ml, 10.8M) and heated to about 70°C. The aqueous phase was separated and washed twice with toluene (300ml each wash). The combined organic washes were concentrated under reduced pressure and the product 1-butyl-4-piperidinylmethylamine (SB-211156) (127g) was isolated as a pale yellow oil in 85% yield by vacuum distillation (bp 106°C at 20mm Hg approx.).

Claims

- 1. A process for the preparation of 1-butyl-4-piperidinylmethylamine, which process comprises:
- 5 i) the reaction of isonipecotamide and 1-bromobutane to give the N-butyl derivative of isonipecotamide; followed by
 - ii) reduction with LiAlH₄,
 characterised in that the reactions i) and ii) are carried out in toluene as solvent.
- 10 2. A process according to Claim 1 in which the reaction mixture after the reduction is treated with hot sodium hydroxide solution and the mixture is extracted with an organic solvent.
- 3. A process for the preparation of SB 207266, or a pharmaceutically acceptable salt thereof, which process comprises preparing 1-butyl-4-piperidinylmethylamine according to the process of claim 1, followed by coupling with an appropriate indole 3-carboxylic acid derivative, and thereafter as necessary converting the indole and/or substituents, including cyclisation to 3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole.
- 20 4. SB 207266 whenever prepared by the process of Claim 3.

INTERNATIONAL SEARCH REPORT

Inter onal Application No PC3/EP 97/05167

A CLASS IPC 6	ification of subject matter C07D211/26 C07D498/04 //(C07	D498/04,265:00,209:00)	
According	o International Patent Classification(IPC) or to both national classi	fication and IPC	
B. FIELDS	SEARCHED		
Minimum d IPC 6	ocumentation searched (classification system followed by classific C 0 7 D	ation symbols)	
Occuments	tion searched other than minimum documentation to the extent the	t such documents are included in the fields se	arched
Electronic	ata base consulted during the international search (name of data	base and, where practical, search terms used	
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Calegory *	Citation of document, with indication, where appropriate, of the r	elevant passages	Relevant to claim No.
A	WO 93 05038 A (SMITHKLINE BEECH March 1993 cited in the application see description 7 and claims	AM PLC) 18	1-4
A	WO 93 18036 A (SMITHKLINE BEECH September 1993 cited in the application see description 1 and claims	AM PLC) 16	1-4
A	WO 94 08965 A (SMITHKLINE BEECH April 1994 cited in the application see description 1-5 and claims	AM PLC) 28	1-4
P,A	WO 96 38420 A (NISSHIN FLOUR MIL LTD.) 5 December 1996 see page 55, example 38	LING CO.,	1,2
Furti	ner documents are listed in the continuation of box C.	X Patent family members are listed in	таплех.
"A" docume consid "E" earlier diling di "L" docume which i citation "O" docume other n	nt which may throw doubts on priority claim(s) or s cited to establish the publicationdate of another or other special reason (as specified) nt reterring to an oral disclosure, use, exhibition or	"T" later document published after the inter- or priority date and not in conflict with cited to understand the principle or the invention. "X" document of particular relevance; the ci- cannot be considered novel or cannot involve an inventive step when the do- "Y" document of particular relevance; the ci- cannot be considered to involve an inv- document is combined with one or mo- ments, such combination being obvious in the art. "5" document member of the same patent f	the application but ory underlying the aimed invention be considered to sument is taken alone aimed invention entive step when the se other such docu— s to a person skilled
	ctual completion of theinternational search	Date of mailing of the international sear	ch report
	3 January 1998	22/01/1998 Authorized officer	
	European Patent Office, P.B. 5816 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Chouly, J	

INTERNATIONAL SEARCH REPORT

information on patent family members

Intr Honal Application No PCT/EP 97/05167

INTERNATIONAL SEARCH REPORT

information on patent family members

Inte Vional Application No PCT/EP 97/05167

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9305038 A		WO 9316072 A JP 7503480 T MX 9300616 A NZ 246915 A ZA 9300764 A AU 4081393 A AU 4350493 A EP 0641198 A WO 9324117 A	19-08-93 13-04-95 01-09-93 28-05-96 26-11-93 30-12-93 24-01-94 08-03-95 09-12-93
WO 9318036 A	16-09-93	AP 401 A AU 671102 B CA 2131797 A CN 1078471 A CZ 9402210 A EP 0630376 A FI 944204 A HU 71121 A IL 105003 A JP 7504433 T MX 9301348 A NO 943348 A NZ 249565 A SI 9300114 A SK 107894 A ZA 9301709 A AU 4081393 A CN 1085083 A EP 0641198 A WO 9324117 A JP 7507290 T MX 9302985 A	29-08-95 15-08-96 16-09-93 17-11-93 13-09-95 28-12-94 12-09-94 28-11-95 12-09-96 18-05-95 01-09-93 09-11-94 27-07-97 31-12-93 12-04-95 18-01-94 30-12-93 13-04-94 08-03-95 09-12-93 10-08-95 01-11-93
WO 9408965 A	28-04-94	EP 0664794 A JP 8502283 T	02-08-95 12-03-96
WO 9638420 A	05-12-96	NONE	